

1 **The unexpected discovery of *Brucella abortus* Buck 19 vaccine in goats from**
2 **Ecuador underlines the importance of biosecurity measures**

3
4 Jorge Ron-Román^{1,2,3,4}, Dirk Berkvens², Daniela Barzallo-Rivadeneira¹, Alexandra
5 Angulo-Cruz¹, Pablo González-Andrade¹, Elizabeth Minda-Aluisa¹, Washington
6 Benítez-Ortíz^{1,5}, Jef Brandt², Richar Rodríguez-Hidalgo¹, Claude Saegerman^{3*}

7
8 ¹ Instituto de Investigación en Salud Pública y Zoonosis, Universidad Central del
9 Ecuador, Quito Ecuador.

10 ² Department of Biomedical Sciences, Institute of Tropical Medicine, Antwerp - Belgium.

11 ³ Research Unit of Epidemiology and Risk analysis applied to Veterinary Sciences
12 (UREAR-ULg), Fundamental and Applied Research for Animal and Health (FARAH)
13 Center, Faculty of Veterinary Medicine, University of Liege, Belgium.

14 ⁴ Carrera de Ingeniería Agropecuaria, Departamento de Ciencias de la Vida y la
15 Agricultura, Universidad de las Fuerzas Armadas (ESPE), Sangolquí, Ecuador.

16 ⁵ Facultad de Medicina Veterinaria y Zootecnia, Universidad Central del Ecuador, Quito,
17 Ecuador.

18
19 **Keywords:** Brucellosis; Goats; Ecuador; Vaccine; Biosecurity.

20
21 *Corresponding author: Research Unit in Epidemiology and Risk Analysis Applied to
22 Veterinary Sciences (UREAR-ULg), Fundamental and Applied Research for Animal and
23 Health (FARAH) Center, Faculty of Veterinary Medicine, University of Liège, B42,
24 Boulevard de Colonster 20, B-4000 Liège, Belgium; e-mail address:
25 claude.saegerman@ulg.ac.be; Tel.: +32-4-366-45-79; Fax: +32-4-366-42-61.

26

27 **Abstract**

28 Very few, mostly old and only preliminary serological studies of brucellosis in goats exist
29 in Ecuador. In order to assess the current epidemiological situation, we performed a cross-
30 sectional serological study in the goat populations of Carchi (n=160 animals), Pichincha
31 (n=224 animals), and Loja provinces (n=2,024 animals). Only two positive serological
32 results (RB negative and SAT-EDTA ≥ 400 IU/ml) were obtained in lactating goats from
33 the same farm in Quito (Pichincha province). Additionally, milk was sampled from 220
34 animals in Pichincha province. The present study indicates a low apparent prevalence in
35 Pichincha province and absence in Carchi and Loja provinces. A total of 25 positive milk
36 ring tests (MRT) were obtained in Pichincha province yielding a prevalence of MRT of
37 11.16 %. Subsequent culture was performed on the positive MRT samples. All results
38 were negative, apart from a single sample, obtained from a serological positive goat in
39 Quito, that was positive for *Brucella abortus* strain 19 (B19). Several hypotheses are
40 forwarded concerning this unexpected result. The most likely hypothesis is the possible
41 accidental use of a needle, previously used for vaccination of cattle with the said vaccine,
42 for the administration of drug treatment to the goat. This hypothesis underlines the
43 necessity of biosecurity measures to prevent this type of accidents.

44

45

46 **Introduction**

47 Brucellosis is a worldwide disease with health and economic impacts (Castro et al.,
48 2005). It is widely distributed in humans and animals, especially in developing countries.
49 Its occurrence is related to the existence of animal reservoirs and high infection rates in
50 livestock, especially in goats and sheep (Corbel, 2006).

51 The main cause of caprine brucellosis is *Brucella melitensis* (biovars 1, 2 and 3)
52 (Godfroid et al., 2010) but some sporadic cases caused by *B. abortus* are documented
53 (e.g., Leal-Klevezas et al., 2000). One or more of the following typically characterize the
54 clinical form of the disease: abortion, retained placenta, orchitis, epididymitis and, more
55 rarely, arthritis together with excretion of the organisms in uterine discharges and milk
56 (OIE, 2016a).

57 Surveillance in goats by indirect diagnostic methods is not a common practice in most
58 countries of South America (PANAFTOSA, 2000), where goat breeding is constrained in
59 its development, because of conditions of overcrowding, poor or non-existent disease
60 control measures and lack of technical assistance, which, together with rudimentary
61 empirical management, permit the transmission of brucellosis (Ortega-Sánchez et al.,
62 2009).

63 Caprine brucellosis due to *Brucella melitensis* is present in Mexico, Peru, Argentina,
64 Paraguay and Bolivia (Aznar et al., 2014; PANAFTOSA, 2000). Until now, there are no
65 reports in Ecuador of isolation and characterization of *Brucella melitensis* in bovines or
66 goats, only molecular findings that demonstrate its presence in samples of lymphatic
67 nodes from goats at the slaughterhouse of Quito (Luna et al., 2016) The total number of
68 goats is estimated between 178,000 (INEC et al., 2002) and 191,000 (OIE, 2016b) of
69 which approximately 43 % (78,000) are found in the canton of Zapotillo in Loja province.

70 The marketing of goat milk in different parts of the Metropolitan District of Quito
71 (two million inhabitants) has become a common activity and forms the basic income of
72 several families engaged in this business. Ecuadorian law prohibits peddling
73 unpasteurized milk, and although vendors work without government regulation, they try
74 as much as possible to maintain minimum health standards, such as collecting animal
75 droppings, washing the udder and selling milk in new and clean bottles (El Comercio,
76 2012).

77 The very few serological studies of brucellosis in goats conducted in Ecuador are old
78 and incomplete or preliminary (e.g., Poulsen et al., 2014). In order to determine the
79 seroprevalence of *Brucella* spp. in goats in three selected areas of Ecuador, as well as
80 isolate the causative agent, we conducted a cross-sectional study (serum and milk
81 samples) in Carchi, Pichincha and Loja provinces.

82

83 **Materials and methods**

84 *Selected areas*

85 The selection of three areas for this study is based on the potential risks: Bolivar and
86 Mira cantons of Carchi province (presence of bovine brucellosis in cattle and existence
87 of mixed farms) (Ron-Román et al. unpublished data), the urban and peri-urban
88 Metropolitan District of Quito in Pichincha province (business of milk goats in Quito city
89 and high density of inhabitants) and Zapotillo canton of Loja (high density of goats)
90 provinces (Figure 1).

91

92 *Sampling design*

93 A survey with census sampling at farm level (n=86) and convenience sampling at
94 animal levels (n=2,408) was performed in the three selected areas. In Carchi and

95 Pichincha provinces (small herds), all herds and all animals present in a herd were
96 sampled. In Zapotillo canton of Loja province (large herds), all herds were included and
97 a random selection of 25 % of animals present in a herd was sampled.

98 In Carchi, blood was sampled between December 2012 and February 2013 (n=160
99 goats in 12 herds). In urban and peri-urban Quito (Pichincha province), blood and milk
100 were sampled between December 2009 and April 2010 (n=224 and 220 goats in 12 herds
101 for blood and milk samples, respectively). In Zapotillo canton of Loja province, blood
102 were sampled in July 2011 (n=2,024 goats in 62 herds). The milk samples were collected
103 only in Quito, area with positive results to serology, to perform the isolation and
104 characterization of the pathogen.

105

106 ***Samples***

107 The goats sampled belonged to native, Nubian and Anglo-Nubian breeds. Jugular vein
108 blood was sampled in vacutainer tubes (10 ml). Each sample was centrifuged; the serum
109 was identified, analysed, and stored at -20 °C. In addition, 100 ml of milk was collected
110 from each lactating goat sampled in peri-urban Quito. All milk samples were identified,
111 stored in a cool box until analysis at the Instituto de Investigación en Salud Pública y
112 Zoonosis (CIZ, Central University of Ecuador).

113

114 ***Blood and milk analysis***

115 Serum samples were analysed for the presence of antibodies against *Brucella* spp.
116 using two diagnostic tests: slide agglutination test with Rose Bengal (RB) and the serum
117 agglutination tube test with EDTA (SAT-EDTA). These tests were performed as
118 previously described (Alton et al., 1988; OIE, 2016a). The modified MRT test as
119 described by Mancera and Ontiveros (2001) for diagnose of brucellosis in goats, was

120 performed as a complementary test on the milk samples. The modification consisted in
121 the addition of 0.3ml of a NaCl solution [25%] and 0.1ml of corn oil to each milk sample
122 (1ml). Afterwards, the samples were incubated at 37°C for 2 hours.

123

124 ***Isolation and identification of Brucella spp.***

125 Milk samples from SAT-EDTA positive (n=2) and MRT positive animals (n=23)
126 were centrifuged at 2,000 g for 15 minutes. The supernatant (cream) and sediment were
127 grown in selective Farrell medium (Columbia Agar Base [Oxoid CM0331] with 5 %
128 decomplexed horse serum [GIBCO Ref-16050-130] and *Brucella* selective
129 supplement [OXOID SR0083A]) for the isolation of *Brucella* spp.

130 Replicated colonies with BASE medium (Columbia Agar Base with 5 %
131 decomplexed horse serum) were identified and classified by means of: macroscopic
132 and microscopic observation, Gram staining and oxidase [DIFCO-BBL Ref: 261181],
133 catalase and urease tests. The procedures were performed as previously described (Alton
134 et al., 1988; Godfroid and Boelaert, 1995).

135

136 ***Identification and molecular characterization of Brucella spp.***

137 Once identified by biochemical tests, the *Brucella* colonies were analysed
138 molecularly by three different PCR tests: the IS6501 PCR or PCR-IS711 (primers: IS6501
139 3': 5'-gat-aga-agg--gct-gaa ctt tgc-gga-c-3' / IS6501 5': 5'-acg-ccg-gtg-tat-ggg-aaa-ggc-
140 ttt-t-3') for genus identification, AMOS PCR (Primers: *B. abortus*-specific: gac-gaa-cgg-
141 aat-ttt-tcc-aat-ccc; *B. melitensis*-specific: aaa-tcg-cgt-cct-tgc-tgg-tct-ga; *B. ovis*-specific:
142 cgg-gtt-ctg-gca-cca-tcg-tcg; *B. suis*-specific: cgc-cgg-ttt-tct-gaa-ggt-tca-gg; IS711-
143 specific: tgc-cga-tca-ctt-aag-ggc-ctt-cat) (Bricker and Halling, 1994) for species
144 determination and modified AMOS PCR (Primers: RB51/2308: ccc-cgg-aag-ata-tgc-ttc-

145 gat-cc; eri primer 1: gcg-ccg-cga-aga-act-tat-caa; eri primer 2: cgc-cat-gtt-agc-ggc-ggt-
146 ga) (Bricker and Halling, 1995) for the differentiation between vaccine strains and field
147 strains.

148

149 *Statistical analysis*

150 The seroprevalence was estimated with a Binomial exact distribution and computed in
151 Stata/MP 14.1 (StataCorp, 2015).

152

153 **Results**

154 No serological RB test showed the presence of antibodies in any of the animals tested
155 but some animals originating from Pichincha province (see below) tested positive for the
156 SAT-EDTA.

157 The study demonstrated the absence of antibodies to *Brucella* spp in Bolivar and Mira
158 cantons of Carchi province (Number of animals tested [Nt]=160; seroprevalence of 0 %
159 with 95 % confidence interval [CI]:0-1.85 %) and Zapotillo canton of Loja province
160 (Nt=2,024; seroprevalence of 0 % with 95 % CI=0-0.15 %). The seroprevalence of
161 brucellosis in the district of Quito in Pichincha province was quite low (Nt=224;
162 seroprevalence of 0.89 % with 95 % CI=0.11-3.19 %).

163 Of the 220 MRT that were performed in Pichincha province, 25 were positive (milk
164 prevalence of 11.16 % with 95 % CI=7.35-16.03 %). Only two goats (out of 47 originating
165 from the same farm in the Tiwinsa sector, urban Quito) were positive in SAT-EDTA (high
166 antibody titres) and in MRT (Table 1). From the two seropositive and lactating goats from
167 Quito urban area, one *Brucella* was isolated on milk. This strain was future characterized
168 and identified as *Brucella abortus* strain 19. The results of the microbiological
169 characterization are in Table 2. A fragment of 498 bp, specific for *Brucella abortus*

170 biotypes 1, 2 or 4, according to Bricker and Halling, (1994), is shown in Figure 2. In
171 Figure 3, the absence of the 364 bp fragment (tandem *IS711*) and the *eri* fragment of 178
172 bp, demonstrate that the strain found in the goat is the B19 vaccine strain (Bricker and
173 Halling, 1995). A further 23 lactating goats that were positive in MRT were negative in
174 culture.

175

176 **Discussion**

177 Brucellosis is a contagious infectious disease, caused by bacteria of the genus
178 *Brucella* spp., which affects both human and several animal species. Caprine brucellosis
179 is mainly due to *B. melitensis* (Godfroid et al., 2010) and some cases of *B. abortus* was
180 previously published (e.g., Leal-Klevezas et al., 2000). The pathogenicity in humans for
181 these two species of *Brucella* is high (Godfroid et al., 2010; Saegerman et al., 2010).

182 The use of SAT-EDTA, RB and MRT was previously evaluated for the diagnosis of
183 caprine brucellosis (Falade, 1978). There was a good correlation between SAT-EDTA
184 and RB when both tests were negative but RB failed to detect 80% of sera above 50 IU/ml
185 in SAT-EDTA. Also, owing to the relatively poor milking potential of the goat and the
186 false positive results with MRT, it was concluded that the SAT-EDTA offers a better
187 serological diagnostic tool for caprine brucellosis. This study is in line with this previous
188 information. Unfortunately, studies reporting serological test results in goats should be
189 interpreted with caution, as most of the data have been obtained without isolation of
190 *Brucella* (Mancera and Ontiveros, 2001).

191 Several preliminary results are available in some Faculties of Veterinary Medicine in
192 Ecuador. In Guayas province (west central part of Ecuador), 33 % of 800 individual milk
193 samples were positive to MRT in 1970 but with no isolation of *Brucella* (Albornoz, 1970).
194 Three other serological studies with Huddleson agglutination test in Macará (Granda,

195 1972), Loja (Tapia, 1998) and Azuay (Sánchez, 1997) provinces indicated a zero or very
196 low seroprevalence.

197 The present study indicates a low prevalence in Pichincha province and absence in
198 Carchi and Loja provinces.

199 The discovery of the *B. abortus* strain 19 (B19) in milk from a goat with a positive
200 serology result (SAW-EDTA: 3,200 IU/ml; high IgM level) was unexpected. Several
201 hypotheses can be postulated. The first hypothesis is the improper use of brucellosis B19
202 vaccine in goats in addition to its advised use in cattle. The brucellosis vaccine of choice
203 for goats is Rev 1 and, as recommended, B19 is only mandatory in cattle in Ecuador and
204 common in Pichincha province. The second hypothesis is a use of a needle, which was
205 previously used for B19 vaccination in cattle, for the administration of a drug to goats.

206 Goats and other species present in a herd are commonly treated by drug injection with the
207 same needle. The second serologically positive goat comes from the same herd, which
208 may form an indication of possible serial use of the same needle. The third hypothesis is
209 the consumption of milk by goats originating from B19 vaccinated cattle. Positive
210 microbiological cultures were obtained during a period of three years from the milk of
211 cows vaccinated with B19 (Meyer and Nelson, 1969), as well as in colostrum (Corner
212 and Alton, 1981). Seropositive titres were observed for a period of one year after B19
213 vaccination of cows (Manthei, 1952). A study of oral vaccination with B19 showed the
214 need of a large dose (500 billion cells) and all serological test were negative in heifers 82
215 days after vaccination (Nicoletti and Milward, 1983). Despite the fact that it cannot be
216 excluded, this hypothesis is deemed unrealistic. The fourth hypothesis is the excretion of
217 B19 in the environment by vaccinated bovines and the use of a same pasture by goats.
218 The intermittent excretion of B19 strain was detected by PCR until 9 years in vaccinated
219 cattle mainly in urine and also in milk samples, which confirmed its multiplication and

220 persistence (Pacheco et al., 2012). However, in this study cultures were always negative.
221 For identical reasons (large dose needed and short period of positivity in serological tests)
222 this hypothesis also appears improbable. In conclusion, the second hypothesis is retained
223 as the most likely.

224

225 **Conclusion**

226 The study demonstrated the absence of antibodies to *Brucella* spp in Bolivar and Mira
227 cantons of Carchi province and Zapotillo canton of Loja province, the principal goat
228 producing canton. Isolation of *Brucella abortus* strain 19 in a goat in Quito district
229 demonstrates the possible cross-infection from vaccinated cattle (B19 vaccination is
230 common here), probably through the accidental use of a needle previously used for
231 vaccination of cattle with B19 vaccine. This finding highlights the necessity of stringent
232 biosecurity measures and quality control of vaccination campaigns.

233

234 **Acknowledgments** This research was funded by the International Centre for Zoonoses,
235 Central University of Ecuador, Quito, Ecuador; the Institute of Tropical Medicine,
236 Antwerp, Belgium and the Research Unit of Epidemiology and Risk analysis applied to
237 Veterinary Sciences, University of Liege, Belgium. The authors thank all farmers who
238 participated in the study.

239

240 **Compliance with ethical standards**

241

242 **Conflict of interest** The authors declare that they have no competing interests.

243

244 **References**

245 Albornoz, G., 1970. Diagnóstico de brucelosis por la prueba de “Ring-test” en la
246 provincia del Guayas a nivel de hacienda. Universidad Estatal de Guayaquil.

247 Alton, G., Jones, L., Angus, R., Verger, J., 1988. Techniques for the brucellosis
248 laboratory, 1st Ed. ed. Paris.

249 Aznar, M.N., Samartino, L.E., Humblet, M.F., Saegerman, C., 2014. Bovine Brucellosis
250 in Argentina and Bordering Countries: Update. Transbound. Emerg. Dis. 61, 121–
251 133. doi:10.1111/tbed.12018

252 Bricker, B.J., Halling, S.M., 1995. Enhancement of the Brucella AMOS PCR assay for
253 differentiation of *Brucella abortus* vaccine strains S19 and RB51. J. Clin. Microbiol.
254 33, 1640–2.

255 Bricker, B.J., Halling, S.M., 1994. Differentiation of *Brucella abortus* bv. 1, 2, and 4,
256 *Brucella melitensis*, *Brucella ovis*, and *Brucella suis* bv. 1 by PCR. J. Clin.
257 Microbiol. 32, 2660–6.

258 Castro, H.A., González, S.R., Prat, M.I., 2005. Brucellosis: una revisión práctica. Acta
259 bioquím. clín. latinoam 39, 203–216.

260 Corbel, M., 2006. Brucellosis in humans and animals. World Health Organization,
261 Geneva Switzerland.

262 Corner, L.A., Alton, G.G., 1981. Persistence of *Brucella abortus* strain 19 infection in
263 adult cattle vaccinated with reduced doses. Res. Vet. Sci. 31, 342–4.

264 El Comercio, 2012. La leche de cabra se vende sin regulaciones [WWW Document]. El
265 Comer. URL [http://www.elcomercio.com/actualidad/quito/leche-de-cabra-se-](http://www.elcomercio.com/actualidad/quito/leche-de-cabra-se-vende.html)
266 [vende.html](http://www.elcomercio.com/actualidad/quito/leche-de-cabra-se-vende.html) (accessed 1.1.12).

267 Falade, S., 1978. A comparison of three serological tests in the diagnosis of caprine
268 brucellosis. Res. Vet. Sci. 24, 376–7.

269 Godfroid, J., Boelaert, F., 1995. Prescriptions pour le diagnostic sérologique de la

270 brucellose. Belgium: CODA-CERVA (Ed.) 47.

271 Godfroid, J., Nielsen, K., Saegerman, C., 2010. Diagnosis of brucellosis in livestock and
272 wildlife. *Croat. Med. J.* 51, 296–305.

273 Granda, B., 1972. Incidencia de brucelosis caprina en el cantón Macará por el método de
274 Huddleson. Universidad Nacional de Loja.

275 INEC, MAG, SICA, 2002. ECUADOR - Agricultural Census 1999/2000 – Main Results
276 [WWW Document]. III Censo Nac. Agropecu. URL
277 http://www.fao.org/fileadmin/templates/ess/ess_test_folder/World_Census_Agricu
278 [lture/Country_info_2000/Reports_2/ECU_SPA_REP_2000.pdf](http://www.fao.org/fileadmin/templates/ess/ess_test_folder/World_Census_Agricu) (accessed 7.20.16).

279 Leal-Klevezas, D.S., Martínez-Vázquez, I.O., García-Cantú, J., López-Merino, A.,
280 Martínez-Soriano, J.P., 2000. Use of polymerase chain reaction to detect *Brucella*
281 *abortus* biovar 1 in infected goats, *Veterinary Microbiology*. doi:10.1016/S0378-
282 1135(00)00200-5

283 Luna, L., Chávez, G., Mejía, L., Barragán, V., Trueba, G., 2016. Molecular Detection of
284 *Brucella* Species in Ecuador. *Intern J Appl Res Vet Med* 14, 185–189.

285 Mancera, A., Ontiveros, M., 2001. Prueba de anillo en leche o anillo de Bang para el
286 diagnóstico de brucelosis en bovinos, in: Díaz, E., Hernández, L., Valero, G.,
287 Arellano, B. (Eds.), *Diagnóstico de Brucelosis Animal*. México, pp. 79–83.

288 Manthei, C.A., 1952. Evaluation of vaccinal methods and doses of *brucella abortus* strain
289 19. *Proc. 56th Annu. Meet. Livest. Sanit. Assoc.* 115–125.

290 Meyer, M.E., Nelson, C.J., 1969. Persistence of *Brucella abortus*, strain 19 infection in
291 immunized cattle., in: *Proceedings, Annual Meeting of the United States Animal*
292 *Health Association*. p. 159.

293 Nicoletti, P., Milward, F.W., 1983. Protection by oral administration of *brucella abortus*
294 strain 19 against an oral challenge exposure with a pathogenic strain of *Brucella*.

295 Am. J. Vet. Res. 44, 1641–3.

296 OIE, 2016a. CHAPTER 2.1.4 Brucellosis (*Brucella abortus*, *B. mellitensis* and *B. suis*)
297 [WWW Document]. OIE. URL
298 http://www.oie.int/fileadmin/Home/esp/Health_standards/tahm/2.01.04_BRUCEL
299 [LOSIS.pdf](#) (accessed 7.20.16).

300 OIE, 2016b. OIE World Animal Health Information System [WWW Document]. WAHIS
301 Interface. URL
302 http://www.oie.int/wahis_2/public/wahidwild.php/Countryinformation/Animalsitua
303 [tion](#) (accessed 7.20.16).

304 Ortega-Sánchez, J.L., Martínez-Romero, A., García-Luján, C., Rodríguez-Martínez, R.,
305 2009. Seroprevalencia de brucelosis caprina en el municipio de Tlahualilo, Durango.
306 México. REDVET. Rev. Electrónica Vet. 10.

307 Pacheco, W.A., Genovez, M.E., Pozzi, C.R., Silva, L.M.P., Azevedo, S.S., Did, C.C.,
308 Piatti, R.M., Pinheiro, E.S., Castro, V., Miyashiro, S., Gambarini, M.L., 2012.
309 Excretion of *Brucella abortus* vaccine B19 strain during a reproductive cycle in
310 dairy cows. Braz. J. Microbiol. 43, 594–601. doi:10.1590/S1517-
311 83822012000200022

312 PANAFTOSA, 2000. Brucellosis y Tuberculosis, situación de los programas en las
313 Américas (No. 1). Rio de Janeiro, Brasil.

314 Poulsen, K.P., Hutchins, F.T., McNulty, C.M., Tremblay, M., Zabala, C., Barragan, V.,
315 Lopez, L., Trueba, G., Bethel, J.W., 2014. Brucellosis in dairy cattle and goats in
316 northern Ecuador. Am. J. Trop. Med. Hyg. 90, 712–5. doi:10.4269/ajtmh.13-0362

317 Saegerman, C., Berkvens, D., Godfroid, J., Walravens, K., 2010. Bovine brucellosis, in:
318 Lefèvre, P., Blancou, J., Chermette, R., Uilenberg, G. (Eds.), Infectious and Parasitic
319 Disease of Livestock. Lovoisier, France, pp. 991–1021.

320 Sánchez, P., 1997. Diagnóstico de brucelosis caprina, en el Cantón Santa Isabel, mediante
321 el método de aglutinación en placa, año 1996. Universidad de Cuenca.

322 StataCorp, 2015. Stata: Release 14. Statistical Software. College Station, TX: StataCorp
323 LP.

324 Tapia, N., 1998. Prevalencia de brucelosis caprina en el área “Centro Laja.” Universidad
325 Nacional de Loja.

326

327

328 CAPTIONS TO ILLUSTRATIONS

329

330 **Figure 1:** Goat population per Canton and localization of the study areas (INEC et al.,
331 2002)

332 Legend: **[A]**, Bolivar and Mira cantons of Carchi province (presence of bovine brucellosis
333 in cattle and existence of mixed farms); **[B]**, urban and peri-urban Metropolitan District
334 of Quito in Pichincha province (business of milk goats in Quito city and high density of
335 inhabitants); **[C]**, Zapotillo canton of Loja province (high density of goats).

336

337 **Figure 2:** PCR amplification products from *Brucella* strains tested by the conventional
338 AMOS assay

339

340 Legend: MP: Molecular weight marker; B1, B2, B3 and B4: Samples of *Brucella* strains
341 by bovines; C1: Samples of *Brucella* strains by caprine (amplification of IS711 which is
342 specific for *B. abortus* biovars 1, 2 or 4 [498 bp]); C-: negative control; C+: positive
343 control of *B. abortus* biovar 1.

344

345 **Figure 3:** PCR amplification products from *B. abortus* strains tested by the modified
346 AMOS assay.

347

348 Legend: MP: Molecular weight marker; B1, B2, B3 and B4: Samples of *B. abortus* strains
349 by bovines; C1: Samples of *Brucella* strains by caprine (absence of amplification of
350 tandem *IS711* [364 bp] and *eri* locus [178 bp]); C-: negative control; C+: positive control
351 of *B. abortus* biovar 1.

352

353

354 **Table 1.** Serology, culture and polymerase chain reaction (PCR) results of two SAT

355 EDTA positive goats

356

Sample N°	Herd Code	Province	Canton	Method of diagnostic						
				RB	SAT-EDTA	MRT	Isolation	PCR IS711	AMOS PCR	mAMOS PCR
178	Tiw 3	Pichinch a	Quito	-	400 IUA	+	-	-	-	-
184	Tiw 3	Pichinch a	Quito	-	3200 IUA	+	+	+	+	+

357

358 Legend: **RB**, Rose Bengal test; **SAT – EDTA**, Serum agglutination test with EDTA;

359 **MRT**, Milk Ring Test **IUA**, International Units of Agglutination **PCR-IS711**,

360 Polymerase chain reaction with insertion 711; **AMOS PCR**, Abortus, Melitensis, Ovis

361 and Suis; **mAMOS PCR**, AMOS modified (PCR for the differentiation of vaccine strains

362 from field strains).

363


364 **Table 2.** Characterization of the caprine *Brucella* spp. isolate

Bacteriological sample code	Catalase	Oxidase	Urease activity	CO ₂ requirement	H ₂ S production	Growth on colorants				Agglutination with serum	
						Thionin 20 µg	Thionin 10 µg	Basic Fuschin 20 µg	Safranin 100 µg	anti A	anti M
Ec-CIZ-Cap-1	+	+++	+(48 hr)	-(48 hr)	+++ (24 hr)	-	-	+	+	+	-
B2*	+	+	+	+	+	-	-	-	-	+	-
B9**	+	+	+	-	+	+	+	+	+	-	+
B1***	+	+	+	+ ^a	+	-	-	+	+	+	-


365

366 Legend: EC-CIZ-Cap-1 is the caprine *Brucella* isolate; * control *Brucella abortus* biovar 2; ** control *Brucella abortus* biovar 9; *** control

367 *Brucella abortus* biovar 1; ^a positive for most strains.




371 **Fig. 2**



372
373

374 **Fig. 3**



375
376