

Smart Point Clouds for information modelling: application in Cultural Heritage

R. Billen, F. Poux & M. Van Ruymbeke

Geomatics Unit, University of Liège



Introduction



About Geomatics Unit


Teaching

- GIS – Mapping
- Surveying – Laserscanning
- Geodesy – GNSS
- Remote Sensing - Photogrammetry


About Geomatics Unit

RB's Research Group


- SI Theory



- 3D GIS



- 3D Reality Capture / 3D Data processing





About Geomatics Unit

RB's Research Group

- SI Theory

- 3D GIS


- 3D Reality Capture / 3D Data processing



About Geomatics Unit

RB's Research Group

- SI Theory
- 3D GIS
- 3D Reality Capture / 3D Data processing



Digital (Built) Cultural Heritage

For Tourism : Video / VR and AR

<https://youtu.be/71vbCPE2w1o>

Digital (Built) Cultural Heritage

For fun ... multi-scale experience ... Creating new space

<https://youtu.be/QNRoYLWBjw8>

Digital (Built) Cultural Heritage

CH information system : CH researches, CH building management
... dealing with all the complexity of CH information

<https://youtu.be/bWHC0kwEtx8>

CH Information system R&D issues


- How to manage multiple needs?
- How to deal with multiple data sources?
- Which data models should be used?
- Do they encompass the whole CH information complexity?
- Etc.



Our researches in Digital (Built) Cultural Heritage

- SI Theory
- 3D GIS
- 3D Reality Capture / 3D Data processing


Smart Point Cloud



Our researches in Digital (Built) Cultural Heritage


- SI Theory
- 3D GIS
- 3D Reality Capture / 3D Data processing

Multiple Interpretation Data Model




Smart Point Cloud


A direct path from 3D Perception to Cognitive Decision



Semantics & Knowledge Integration



TODAY
←→
WHAT WE WANT




KNOWLEDGE




DISCOVERY FROM EXTRACTION
plan, based on REASONING, ...

Conceptual SPC Model Overview




Level-0 SPC Meta-Model

GENERALIZED SPC META-MODEL




Level-1: SPC CL Meta-Model




CONNECTION-LAYER META-MODEL

L1-1




L1-2




Level-2 SPC Domain Adaptation

CONNECTION-LAYER META-MODEL L1-2



DOMAIN META-MODEL



SPC application to CH



Device knowledge



TLS DATA



DENSE-IMAGE MATCHING DATA




Fusion / NURBS



Colour point cloud

Device knowledge


- Amplitude of the spatial error between TLS and NURBS
- TLS signal is influenced by the material, the error issued from the material reflectance can be used as a semantic information to help the classification process.



Analytical knowledge




Colour point cloud



Segmented point cloud

Type	Point Features	Range	Explanation
Sensor desc.	X, Y, Z	Bounding-box	Limits the study of points to the zone of interest
	R, G, B ¹	Material Colour	Limited to the colour range that domain knowledge specifies
	I		Clear noise and weight low intensity values for signal representativity
Shape desc.	RANSAC ²	-	Used to provide estimator of planarity
Local desc.	N _x , N _y , N _z ³	[-1, 1]	Normalized normal to provide insight on point and object orientation
	Density ⁴	-	Used to provide insights on noise level and point grouping into one object
	Curvature	[0, 1]	Used to provide insight for edge extraction and break lines
	KB ⁵ Distance map		Amplitude of the spatial error between the raw measurements and the final dataset
Structure desc. ⁶	Voxels	-	Used to infer initial spatial connectivity


Domain knowledge




Domain knowledge












Colour point cloud



Segmented point cloud



Classified point cloud


Sample	Available knowledge			
	Surface (in cm ²)	Approx. Geometry	Reflectance (at 1.55 μm)	Date
 Gold	~ 1 cm ²	~ square	H. Reflective	XIX
 Gold	< 1 cm ²	irregular	~ Mat	<
 Faience	~ 20 cm ²	tear, diamond	Reflective	XIX
 Silver	~ 1 cm ²	~ square	H. Reflective	XIX
 C. glass	~ 1 cm ²	irregular	~ Reflective	<
 C. glass	~ 1 cm ²	irregular	~ Absorbent	<
 C. glass	~ 1 cm ²	irregular	Reflective	<
 C. glass	~ 1 cm ²	irregular	Reflective	<
 C. glass	~ 1 cm ²	irregular	~ Mat	<

Reasoning

Language	RDF Triple Store	Effect
SPARQL	<pre> PREFIX rdf:<http://www.w3.org/1999/02/22-rdf-syntax-ns#> PREFIX npt: <http://www.geo.ulg.ac.be/nyspoux/> SELECT ?ind WHERE { ?ind rdf:type npt:AltoMedievalTessera } ORDER BY ?ind </pre>	Return all alto-medieval tesserae (regarding initial data input)
SQL	<pre> SELECT name, area FROM worldObject WHERE ST_3DIntersects(geomWo::geometry, polygonZ::geometry); </pre>	Return all tesserae which are comprised in the region defined by a selection polygon and gives their area
SPARQL & SQL	<pre> SELECT geomWo FROM worldObject WHERE ST_3DIntersects(geomWo::geometry, polygon2Z::geometry) AND area > 0,0001; PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#> PREFIX npt: <http://www.geo.ulg.ac.be/nyspoux/> SELECT ?ind WHERE { ?ind rdf:type npt: XIXCentTessera } ORDER BY ?ind </pre>	Return all renovated tesserae in the region 2 where the area is superior to 1 cm ²

Semantic integration by user / training platform

Multiple Interpretation Data Model



CIDOC-CRM MDIM extension



What is the MIDM?

Multiple Interpretation Data Model

We aimed at developing a model handling ...

- Complex nature of CH information and its inherent imperfection
- Events
- Multiple interpretation

Peculiarities of archaeological Data




What is the MIDM?

Multiple Interpretation Data Model


We aimed at developing a model handling ...

- Complex nature of CH information and its inherent imperfection
- Events
- Multiple interpretation

What is the MIDM?



What is the MIDM?



MIDM - Example

- Record of two Episodes (version) about the Church of Theux

EP-V1
Church with Tower 1175

EP-V2
Gothic choir 1520


What is the MIDM?

Multiple Interpretation Data Model

We aimed at developing a model handling ...


- Complex nature of CH information and its inherent imperfection
- Events
- Multiple interpretation

What is the MIDM?




MIDM - Example

Record of one Interpretative
Sequence by author 1



MIDM - Example

Life Map of the Church of Theux




MIDM - Example

EP-V1
Church with Tower 1175

EP-V2
Gothic choir 1520


MIDM - Example

Record of new Episodes (version and event) about the Church of Theux




MIDM - Example

Record of one Interpretative Sequence by author 2




MIDM - Example

Record of a second Interpretative Sequence by author 2




MIDM - Example

Record of new Episodes (version and event) about the Church of Theux




MIDM - Example

Record of one Interpretative Sequence by author 3



MIDM - Example

New Life Map of the Church of Theux



MIDM Future


- MIDM first version has been mapped to CityGML
 - Chaturvedi K., Smyth C.S., Gesquière G., Kutzner T., Kolbe T.H. (2017) Managing Versions and History Within Semantic 3D City Models for the Next Generation of CityGML. In: Abdul-Rahman A. (eds) Advances in 3D Geoinformation. Lecture Notes in Geoinformation and Cartography. Springer, Cham



- MIDM last version has been mapped to CIDOC CRM



Is MIDM just for CH and history?



Conclusions

- SPC is a new way to use point cloud as structural spatial elements carrying semantic information
- SPC is obviously not limited to CH application... can be plugged to other domains

- MIDM is a new way to encompass complexity of CH information and variability of experts' interpretations
- MIDM not only model the past but could be used to model the uncertainty of the future

These papers can interest you (SPC):

- Poux, F., Hallot, P., Neuville, R., Billen, R., 2016. SMART POINT CLOUD: DEFINITION AND REMAINING CHALLENGES. ISPRS Ann. Photogramm. Remote Sens. Spat. Inf. Sci. IV-2/W1, 119–127. doi:10.5194/isprs-annals-IV-2-W1-119-2016
- Poux, F., Neuville, R., Billen, R., 2017a. POINT CLOUD CLASSIFICATION OF TESSERAE FROM TERRESTRIAL LASER DATA COMBINED WITH DENSE IMAGE MATCHING FOR ARCHAEOLOGICAL INFORMATION EXTRACTION. ISPRS Ann. Photogramm. Remote Sens. Spat. Inf. Sci. IV-2/W2, 203–211. doi:10.5194/isprs-annals-IV-2-W2-203-2017
- Poux, F., Neuville, R., Hallot, P., Billen, R., 2017b. MODEL FOR REASONING FROM SEMANTICALLY RICH POINT CLOUD DATA. ISPRS Ann. Photogramm. Remote Sens. Spat. Inf. Sci. in press.
- Poux, F., Neuville, R., Wersch, L. Van, Nys, G.-A., Billen, R., 2017c. 3D Point Clouds in Archaeology: Advances in Acquisition, Processing and Knowledge Integration Applied to Quasi-Planar Objects. Geosci. 2017, Vol. 7, Page 96 7, 96. doi:10.3390/GEOSCIENCES7040096

These papers can interest you (MIDM):

- Pfeiffer, M., Carré, C., Delfosse, V., Hallot, P., & Billen, R. (2013). Virtual Leodium: from an historical 3D city scale model to an archaeological information system. *ISPRS Annals–Volume II-5/W1, 2013*.
- Van Ruymbeke, M., Carré, C., Delfosse, V., Pfeiffer, M., & Billen, R. (2015). Towards an Archaeological Information System: improving the core data model. In *CAA 2014 21st century Archaeology: Concepts methods and tools: Proceedings of the 42nd Annual Conference on Computer Applications and Quantitative Methods in Archaeology* (pp. 245-253). Archaeopress.
- Van Ruymbeke, M, Hallot, P. & Billen, R. (2017). Enhancing CIDOC - CRM and compatible models with the concept of multiple interpretation. *ISPRS Annals of the Photogrammetry, Remote Sensing and Spatial Information Sciences, Volume IV-2/W2 , 287-294*.
- Van Ruymbeke, M, Hallot, P. & Billen, R. (Forthcoming) IMPLEMENTATION OF MULTIPLE INTERPRETATION CONCEPT IN CIDOC-CRM AND COMPATIBLE MODELS, *Virtual Archaeology Review*

Thank you !

fpoux@uliege.be, rbillen@uliege.be

geomatics.uliege.be (soon ;-))

geomatics.ulg.ac.be (temporary)

